No.



9000123

TO ARE TO WHOM THESE PRESENTS SHARE COME:

Golden's Foundation Seeds, Inc.

Withereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-CLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT TY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN

'LH162'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affired at the City of Washington, D. C. 29th day of

the year of our Lord one thousand nine

hundred and ninety-one.

Plant Variety Protection Office Spricultural Marketing Service

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, OIRM, Room 404-W, Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055). Washington, 20250.

U.S. DEPARTMENT OF A AGRICULTURAL MARKET	GRICULTURE ING SERVICE		Application is required in order to
APPLICATION FOR PLANT VARIETY (Instructions on I		CERTIFICATE	determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).
NAME OF APPLICANT(S) (as it is to appear on the Certificate)		2. TEMPORARY DESIGNATION OR EXPERIMENTAL NO.	3. VARIETY NAME
Holden's Foundation Seeds, Inc.	1	Ex1608	LH162
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)		5. PHONE (Include area code)	FOR OFFICIAL USE ONLY
201 N. Maplewood Avenue			PVPO NUMBER
P.O. Box 839		319-668-1100	9000123
Williamsburg, Iowa 52361			F Date
6. GENUS AND SPECIES NAME	7. FAMILY NAME (Botania	L	1 71/1000 23,1990
Zea mays	Gramineae		N
8. CROP KIND NAME (Common Name)	9.	DATE OF DETERMINATION	F Filing and Examination Fee:
Corn, Field	i	November 1989	S Date
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGAN	NZATION (Corporation, part	nership, association, etc.)	R Than. 23,1990
Corporation			C Certificate Fee:
11. IF INCORPORATED, GIVE STATE OF INCORPORATION	12. DA	TE OF INCORPORATION	E \$ 250.
Iowa		1968	5 Feb. 19, 1991
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO	SERVE IN THIS APPLICATION	ON AND RECEIVE ALL PAPERS	
Mr. Mark Armstrong			
P.O. Box 839			010 660 1100
Williamsburg, Iowa 52361		PHONE (Include area cod	319-668-1100
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Folician Exhibit A, Origin and Breeding History of the Variety. b. X Exhibit B, Novelty Statement. c. Exhibit C, Objective Description of Variety. d. X Exhibit D, Additional Description of Variety. e. X Exhibit E, Statement of the Basis of Applicant's Ownership Seed Sample (2,500 viable untreated seeds). Date Seed G. X Filing and Examination Fee (\$2,150) made payable to "Trees."	p. Sample mailed to Plant V	fariety Protection Office $3/2//$	<u>96</u>
15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SOL Protection Act.)	D BY VARIETY NAME ONL	AS A CLASS OF CERTIFIED SEED? (Se	e section 83(a) of the Plant Variety
YES (If "YES," answer items 16 and 17 belong 16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS T		O," skip to item 18 below) DITEM 16, WHICH CLASSES OF PRODU	OTION REVOND RREEDED CEEDS
NUMBER OF GENERATIONS?			
L YES L NO	∐ FOL	NDATION REGIST	ERED CERTIFIED
18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION OF THE VAR	RIETY IN THE U.S.?		
☐ YES (If "YES," Ihrough ☐ Plant Variety Protection Act ▼ NO	Patent Act. Give da	e:)	
19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR MA	ARKETED IN THE U.S. OR (THER COUNTRIES?	
YES (If "YES," give names of countries and dates) NO			
20. The applicant(s) declare(s) that a viable sample of basic see request in accordance with such regulations as may be appli	eds of this variety will	be furnished with the application	on and will be replenished upon
The undersigned applicant(s) is (are) the owner(s) of this uniform, and stable as required in section 41, and is entitled	sexually reproduced	novel plant variety, and believe	e(s) that the variety is distinct,
Applicant(s) is (are) informed that false representation here			tant fatiety i toechon net.
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR	TITLE	DATE
(/ molel Tfolder	Presid	lent	3/2//90
SIGNATURE OF APPLICANT [Owner(s)]	CAPACITY OR	TITLE	DATE

Origin and Breeding History of the Inbred

Exhibit A

LH162 was developed from a cross ND246 x Early Mo17 Pollen Composite. The seed from this cross was selfed and the pedigree system of plant breeding was used to develop LH162. ND246 is a public corn inbred line developed and released by North Dakota State University. The Early Mo17 Composite is a population of early maturing Mo17 type plants developed by and the property of Holden's Foundation Seeds, Inc. The earliest maturing plants were selected from the Early Mo17 Composite population and a bulk pollen sample was gathered and used to cross to ND246. On the following pages are a summary and description of the development of LH162. Also included are copies of pages from Holden's Foundation Seeds, Inc. nursery books. The rows associated with the development of LH162 have been highlighted.

Attached is a statement from the originating plant breeder, Scott Bergemann, Holden's Foundation Seeds, Inc., stating that the line, LH62, is stable, uniform and free of variance within the population.

Origin and Breeding History of the Inbred LH162 = Ex1608 = ND246 x Early Mo17

Exhibit A

Row/Field	<u>Pedigree</u>	Location	<u>Year</u>
Home Field	LH162	Iowa	1989
West Elwood Garden	LH162	Iowa	1989
Home Field	LH162	Iowa	1988
2879-2898	LH162	Minnesota	1988
201-210	Ex1608	Minnesota	1987
585	ND246 x Early Mo17	Minnesota	1986
14947	ND246 x Early Mo17	Hawaii	1985-86
8218	ND246 x Early Mo17	Minnesota	1985
21411	ND246 x Early Mo17	Hawaii	1984-85
5566	ND246 x Early Mo17	Minnesota	1984
7017	ND246 x Early Mo17	Hawaii	1983-84
5583	ND246 x Early Mo17	Minnesota	1983
1-469	Early Mo17 Composite	Iowa	1982
15391	ND246	Iowa	1982

Uniformity Statement

Exhibit A

I have observed LH162 for the past three generations it has been increased, 1987 Minnesota nursery rows 201-210, 1988 Minnesota nursery rows 2879-2898, 1988 Iowa production Home Field and 1989 Iowa production Home Field and West Elwood Garden Field. In each of these increases, seeds from the previous generation were planted. The line is very stable and uniform. LH162 is also free of variance from within the population.

Scott Bergemann

Plant Breeder

Novelty Statement

Exhibit B

LH162 most closely resembles the corn inbred line ND246, however, the tassel branches of LH162 are more erect than the tassel branches of ND246. The tassel branches of ND246 are not erect and tend to droop or "flatten out". Photographs of both inbred tassels are attached showing this difference.


The silk color of LH162 is salmon, while the silk color of ND246 is green. A photograph of the exposed silks of LH162 is attached.

LH162 has a pink cob color, whereas the cob of ND246 is red.





Silk color of LH162



FORM GR-470-28 (2-15-74)

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION

EXHIBIT C (Com)

HYATTSVILLE, MARYLAND 20782

OBJECTIVE DESCRIPTION OF VARIETY

CORN (ZEA MAYS)

	NAME OF ADDITIONAL OF ADDITION			
	NAME OF APPLICANT(S)	FOR OFFICIAL USE ONLY		
_	Holden's Foundation Seeds, Inc. ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	9000123		
	201 N. Maplewood Avenue	VARIETY NAME OR TEMPORARY		
	P.O. Box 839	DESIGNATION		
	Williamsburg, Iowa 52361	LH162		
—	Place the appropriate number that describes the varietal character of this variety in the			
	Place a zero in first box (e.s. 0 8 9 or 0 9) when number is either 99 or less or	9 or less.		
	1. TYPE:			
	$2\&\beta$ 1=SWEET 2=DENT 3=FLINT 4=FLOUR 5=P	OP 6 = ORNA. ENTAL		
	2. REGION WHERE BEST ADAPTED IN THE U.S.A.:			
	1 = NORTHWEST 2 = NORTHCENTRAL 3 = NORTHEAST	4 = SOUTHEAST		
	5 - SOUTHCENTIAL 6 - SOUTHWEST / = MOST REGIONS			
		comments" (pg. 3) state how		
		's were calculated)		
	7 1 DAYS FROM EMERGENCE TO 50% OF PLANTS IN SILK 1 2	2 1 HEAT UNITS		
	0 0 DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY	0 0 HEAT UNITS		
				
	0 0 DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 0 0	0 0 HEAT UNITS		
	4. PLANT:			
	1 7 1 CM. HEIGHT (To tassel tip)	8 5 CM. EAR HEIGHT (To base of top ear)		
	[
	1 1 1 CM. LENGTH OF TOP EAR INTERNODE			
	Number of Tillers: Number of Ears Per Stalk			
Number of Fillers: Number of Ears Per Stalk:				
		SLIGHT TWO-EAR TENDENCY		
	3 = STRONG TWO	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 = THREE-EAR TENDENCY		
	Cytoplasm Type:	-EAR TENDENCY 4 = THREE-EAR TENDENCY		
	3 = STRONG TWO	-EAR TENDENCY 4 = THREE-EAR TENDENCY		
	Cytoplasm Type:	-EAR TENDENCY 4 = THREE-EAR TENDENCY		
-	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given):	-EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue	-EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given):	-EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN	-EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue	-EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence:	-EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify) SS EEN (B14) 4 = VERY DARK GREEN (K166)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN	(Specify) SS 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREATER (MF9) 3 = DARK GREATER (MF9	(Specify) SS 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN (HY) Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° 2 1 = LIGHT	(Specify) SS 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREATER (MF9) 3 = DARK GREATER (MF9	(Specify) (Specify) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° 2 1 = LIGHT 3 = HEAVY Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSEN 3 = MANY	(Specify) (Specify) 2S EEN (B14) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26) T (OH51) 2 = FEW (OH56A)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° 2 1 = LIGHT 3 = HEAVY Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSEN 3 = MANY Width: Length:	(Specify) (Specify) 2S EEN (B14) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26) T (OH51) 2 = FEW (OH56A)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN (WF9) 3 = DARK GREEN (WF9) 3 = DARK GREEN (WF9) 3 = MANY (OH7L) Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSEN 3 = MANY Width: Length:	(Specify) (Specify) (Specify) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26) (OH26) (OH51) 2 = FEW (OH56A) (PA11)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° 2 1 = LIGHT 3 = HEAVY Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSEN 3 = MANY Width: Length:	(Specify) (Specify) 2S EEN (B14) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26) T (OH51) 2 = FEW (OH56A)		
	Cytoplasm Type: 1 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 5GY 4/6 Munsell Color Charts for Plant Tissue 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN (WF9) 3 = MANY (OH7L) 2 1 = ABSEN 3 = MANY (OH7L) 2 1 = ABSEN 3 = MANY (OH7L) 2 Length:	(Specify) (Specify) (Specify) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26) (OH26) (OH51) 2 = FEW (OH56A) (PA11)		

Shape Grade (% Rounds)

1 = < 20

2 = 20-40

3 = 40-60

4 = 60-80

4

RM	GR-470-28	Page 2 of 3
6.	TASSEL:	
	1 1 NUMBER OF LATERAL BRANCHES	000123
	Branch Angle from Central Spike: Penduncle Length:	•
	$2 \qquad 1 = \langle 30^{\circ} \qquad 2 = 30 - 40^{\circ} \qquad 3 = \rangle 45^{\circ} \qquad \boxed{0 4} \qquad CM. FROM TOP LEAF TO BASAL BRA$	NCHES
	Pollen Shed:	
	2 1 = LIGHT (WF9) 2 = MEDIUM 3 = HEAVY(KY21)	
	*bleaching tan	
	Anther Color: 1 = YELLOW 2 = PINK 3 = RED 4 = PURPLE 5 = GF Glume Color: 6 = OTHER (Specify) green with brown margin	REEN
	Pollen Restoration for Cytoplasms (o = Not Tested, 1 = Partial, 2 = Good)	
	0 "T" 0 "S" 0 "C" 0 OTHER (Specify Cytoplasm and degrees of restoration)	
7	EAR (Husked Ear Data Except When Stated Otherwise):	
	1 6 CM LENGTH 4 0 MM. MID-POINT 9 0 GM. WEIGHT	
	Kernel Rows:	
	2 1 = INDISTINCT 2 = DISTINCT 1 4 NUMBER	
	1 = STRAIGHT 2 = SLIGHTLY CURVED 3 = SPIRAL	
	Silk Color (Exposed at Silking Stage):	
	1 = GREEN 2 = PINK 3 = SALMON 4 = RED	
	Husk Color:	
	FRESH 1 = LIGHT GREEN 2 = DARK GREEN 3 = PINK	
	6 DRY	
	lusk Extention: (Harvest Stage) Husk Leaf:	
	1 = SHORT (Ears Exposed) 2 = MEDIUM (Barely Covering Ear) 2 3 = LONG (8-10CM Beyond Ear Tip) 1 = SHORT (< 8 CM) 2 = MEDIUM (8-10CM Sevend Ear Tip) 3 = LONG (> 15 CM)	-15 CM)
	4 = VERY LONG (> 10 CM) hank: Position at Dry Husk Stage:	
	1 4 CM LONG 7 NO. OF INTERNODES 1 1 = UPRIGHT 2 = HORIZONTAL	3 = PENDENT
	aper: Drying Time (Unhusked Ear):	
	2	3 = FAST
8.	CERNEL (Dried):	
	ize (From Ear Mid-Point):	
	1 1 MM LONG 0 8 MM. WIDE 0 4 MM. THICK	

5 = > 80

8. KERNEI	_ (Dried) :					-	·
1	Pericarp Color:	1 = COLORLESS 5 = BROWN 8 = VARIEGATED (Descrit	6 = LIGHT F		3 = TAN 7 ⇒ CHERRY R	4 = BRONZE ED	
1	Aleurone Color:	1 = HOMOZYGOUS	2 = SEGF	REGATING (Describe)			
1		2 = PINK 3 = 8 = PALE PURPLE		-		5 = BRONZE	6 = RED
3	Endosperm Color:	1 = WHITE 2 = PA	LE YELLOW	3 = YELLOW	4 = PINK-ORA	NGE 5 = WHI	ГЕ САР.
Endosper	т Туре:						
3		2 = EXTRA SWE CH 6 = HIGH PROTE		3 = NORMAL STA	0 - 1	HIGH AMY L OSE STA OTHER (Specify)	ARCH
2 0	GM. WEIGHT /100	SEEDS (Unsized Sample)					***************************************
9. COB:							
2 9	MM. DIAMETER A	AT MID-POINT					
Strength	:		Со	lor:			
	1 = WEAK	2 = STRONG		1		ED. 4 = BROWN	
ــــــــــــــــــــــــــــــــــــــ			<u> </u>	5 = VARIEGATE	D 6 OT I	HER (Specify)	
10. DISEAS	E RESISTANCE (O	= Not Tested, 1 = Susceptible,	2 = Resistant):				
0	0 STALK ROT (Diplodia)				a)		
0	NORTHERN LEA	F BLIGHT 0 S	OUTHERN LE	AF BLIGHT	0 s	иит	
	SOUTHERN RUST	r	CORN SMUT		0 B/	ACTERIAL WILT	
	BACTERIAL LEA	FBLIGHT 0	MAIZE DWARF	MOSAIC	ST ST	UNT	
0	OTHER (Specify)	[0]					
	PECISTANCT IO =	Not Tested, 1 = Susceptible, 2	P = Posistant):				
	CORNBORER	0 EARWORI		0 SA	PBEET L E	O APHIE)
0	ROOTWORM (Nor	rthern) 0 ROOTWO	RM (Western)			ي ا	
	ROOTWORM (Sou	others (c					
12. VARIET	TES MOST CLOSEL	Y RESEMBLING THAT SUB	MITTED FOR	THE CHARACTERS	GIVEN:		
CHARA	CTER	VARIE	TY	CHARACTE	R	VARIETY	
Maturity Plant Tw		CM105		Kernel Type	-te\	ND246	
Plant Type		ND246 ND246		Quality (Edib Usage	ole)	ND246	
REFERE	NCES: U.S. Department A	griculture. Yearbook 1937.					
		cessing, Products. 1970 Avi F			•		
		N. Beadle, and A.C. Fraser. A ize. 1968. Crop Science Socience		_	.Cornell A.E.S., M	em. 180. 1935.	
		aize Inbred Lines of Ohio. Oh					
		- A System for the Classifica			sis, Ohio State Un	iversity.	

 $\frac{\text{Tmax} + \text{Tmin}}{2}$

GDD =

- 50°F

COMMENTS:

Tmax <u>≤</u>86°F Tmin ≥50°F

Additional Description of the Inbred

Exhibit D

LH162 is taller in plant height and ear height than ND246. Measurements taken during the growing season of 1989 at Williamsburg, Iowa, showed LH162 had an average plant height of 171 centimeters and an average ear height of 85 centimeters. ND246 had an average plant height of 163 centimeters and an average ear height of 69 centimeters.

The silk color of LH162 is salmon, while the silk color of ND246 is green.

The ears of LH162 and ND246 are similar in length, however the diameter of the LH162 ear is greater than the diameter of the ND246 ear (40 cm vs. 35 cm). This difference is due to the greater kernel depth or greater kernel length of LH162. A kernel of LH162 measured at the mid-point of the ear measures 11 mm long, 4 mm thick and 8 mm wide. A kernel of ND246 measured at the mid-point of the ear measures 9 mm long, 4 mm thick and 7 mm wide.

LH162 has a pink cob color, whereas the cob color of ND246 is red.

LH162 performs best when crossed to early B14 lines. In these hybrids, LH162 contributes early flowering, early maturity and good standability. LH162 appears to be a good pollinator and an average female.

- Novelty-Statement

JMS 1/11/91

Exhibit & D addendum

LH162 most closely resembles the corn inbred line ND246, however, the most distinguishing characteristic is the tassel. The tassel of LH162 has a more closed or upright tassel branch angle (1865 than 30°), while the tassel of ND246 is more open and not as upright (30°-40°). The photograph of the two tassels below illustrates this characteristic.

IMS 1/11/91




Exhibit E

STATEMENT OF THE BASIS OF APPLICANT OWNERSHIP

Holden's Foundation Seeds, Inc., Williamsburg, Iowa, is the sole owner and breeder of the LH162 corn inbred line for which it solicits a certificate of protection.